

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

## Intracellular calcium signaling: holding the balance between health and disease.

### This is the author's manuscript

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/117396> since 2015-12-23T12:23:23Z

*Published version:*

DOI:10.2174/092986712804143330

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)



# UNIVERSITÀ DEGLI STUDI DI TORINO

***This is an author version of the contribution published on:***

*Questa è la versione dell'autore dell'opera:*

*[inserire titolo della rivista, volume e fascicolo, anno, DOI]*

*ovvero [autore, volume, editore, anno, pagg.XXXX-XXXX]*

***The definitive version is available at:***

*La versione definitiva è disponibile alla URL:*

*[inserire URL sito editoriale]*

## EDITORIAL

### **Intracellular calcium signaling: holding the balance between health and disease.**

Flexibility and conservation are two hallmarks of intracellular  $\text{Ca}^{2+}$  ( $\text{Ca}_c$ ) signaling. These intriguing features contributed to fix this ion as a prominent chemical regulator in cell physiology [1-5].

Calcium machinery, the entire toolkit of biomolecules generating and decoding  $\text{Ca}_c$  events inside the cell, includes a number of components, some ubiquitous and others specifically expressed by some tissues [3]. Several proteins, often clustered in multimers, compose ion channels that allow  $\text{Ca}^{2+}$  fluxes into the cytosol. They can be activated by mechanical forces, changes in membrane potential, intracellular or extracellular chemical agonists, and depletion of intracellular  $\text{Ca}^{2+}$  stores [3]. Nonetheless, some of them, typically transient receptor potential channels (TRPs), are multifunctional and can be simultaneously modulated through multiple mechanisms [6, 7].

Calcium signaling is tightly coupled to cell metabolism in a reciprocal crosstalk. Consistently, a number of diseases are linked to alterations in  $\text{Ca}_c$  signals, that can act as a cause as well as secondary effects of the pathological progression [8-11]. Furthermore  $\text{Ca}_c$  machinery, that shapes  $\text{Ca}_c$  waves and decodes  $\text{Ca}_c$  'signature', does not work as an isolated system, being integrated with other cell signaling pathways and intracellular messengers. They include phosphoinositides ( $\text{InsP}_3$ ,  $\text{PIP}_2$ ), gasotransmitters ( $\text{NO}$ ,  $\text{H}_2\text{S}$  and  $\text{CO}$ ), bioactive lipids (arachidonic acid, eicosanoids, diacylglycerol), and cyclic nucleotides (cAMP and cGMP) [5, 12-19]. cAMP is released during stimulation with hormones and growth factors. It was the first intracellular messenger discovered in 1971, and, similarly to  $\text{Ca}^{2+}$ , is universal and evolutionary conserved [20]. Recent research has revealed that cAMP-mediated signaling is highly compartmentalized and relies on a complex network of intracellular pathways, whose relative relevance depends on the specific extracellular stimulus. Mutations or genetic polymorphism involving some cAMP-related components are

associated with a variety of human diseases including long QT syndrome, cardiac dysfunctions, familial breast cancer and schizophrenia [20]. The review by A. Hofer focuses on the complex crosstalk between  $\text{Ca}_c$ - and cAMP-dependent networks. It is well known that cAMP microdomains modulate the activity of several  $\text{Ca}^{2+}$  channels, either directly or indirectly, *via* phosphorylation by cAMP-dependent protein kinase, PKA, and cAMP sensor Epac (exchange proteins activated directly by cAMP) [18]. On the other hand, the cAMP pathway is subject to modifications by  $\text{Ca}^{2+}$  and its effectors at many levels. Consistently, the authors recently reported a mechanism of cAMP release dependent on intracellular calcium store depletion, called ‘Store-Operated cAMP Signaling’ or ‘SOcAMPS’ [18-21]. Excessive activation of store-operated cAMP production has been proposed to cause human progressive polycystic liver disease [22].

Specificity of  $\text{Ca}^{2+}$ -dependent biological effects can be achieved through the selective expression and targeting of given  $\text{Ca}^{2+}$  channels and effectors. In addition, a great amount of data point to the relevance of localized intracellular events,  $\text{Ca}_c$  microdomains, due to ‘signalosomes’, organized clusters of  $\text{Ca}^{2+}$  channels, transporters and  $\text{Ca}^{2+}$ -dependent interactors [23, 24]. The review by Ambudkar provides a striking example of  $\text{Ca}_c$  signaling anisotropy: the exocrine function in secretory epithelia. Salivary and pancreatic gland acinar cells secrete fluid, ions, and proteins in a vectorial process that requires the coordinated regulation of ion and water channels and transporters, as well as of signaling molecules and vesicles. Spatially restricted  $\text{Ca}_c$  signals play a critical role in the regulation of this complex phenomenon [25]. The polarized co-localization of both  $\text{Ca}_c$  signaling and secretory components accounts for the physiological vectorial secretion in the exocrine salivary glands and pancreas. Pathological disruption of calcium handling is associated with autoimmune inflammatory diseases such as acute pancreatitis and Sjögren's syndrome [26, 27]. This is a fascinating example that explains how the physiopathological implications of altered  $\text{Ca}_c$  signaling at the single cell level depend on tissutal organization.

Among the major leading causes of death in the world, heart diseases are object of an impressive amount of basic and clinical studies at both cellular and tissutal levels. On the basis of its calcium-dependent excitability, it is not surprising that a number of cardiac dysfunctions in adults can be successfully related to alterations occurring in the  $\text{Ca}_c$  homeostasis of single cardiomyocytes [28, 29]. In this issue, Levi et al. investigate in detail the role of  $\text{Ca}_c$  signals in heart failure and protection from damage [30, 31]. Cardiac development, growth and differentiation are physiologically controlled by the action of several growth factors. It has been recently shown that the exogenous administration of these peptides protects the heart from failure and ischemia/reperfusion (I/R) injury [32]. Moreover, cardioprotection involves the activation of kinases, such as the so-called Reperfusion Injury Salvage Kinase (RISK) pathway, that includes phosphoinositide 3 kinase (PI3K)/Akt and extracellular regulated signal kinases 1 and 2 (Erk1/2). Unfortunately, continuous activation of RISK cascades can result in the loss of their cardioprotective effect, hypertrophic responses and undesired secondary effects.

The authors discuss novel putative protective mechanisms that involve peptides acting through  $\text{Ca}_c$  signaling (i.e. neuregulin, urocortin and natriuretic peptides) and that, as in the case of the Neuregulin-1beta/ErbB pathway, are reaching the clinical trial relevance. Particular attention is focused on SUMOylation, a novel post-translational mechanism acting in cardioprotective processes and cardiac  $\text{Ca}_c$  handling [33].

Intracellular  $\text{Ca}^{2+}$  signals drive morphogenetic processes at tissutal level that, in turn, exert a feedback on single cells, giving rise to a functional loop between different biological scales [34, 35]. This issue includes two reviews focused on the putative multiple roles of calcium in particular biological processes, such as neuronal migration during central nervous system development and endothelial functions during the formation of new blood vessels, angiogenesis.

Lovisolo et al. discuss the role of  $\text{Ca}_c$  signaling on radial and tangential migration of neurons and neuronal precursors during the development of the nervous system [36]. Deciphering the involvement of the different  $\text{Ca}^{2+}$  influx pathways has been a major task for cellular neurobiologists, in which the main limiting factor is the limited availability of reliable and selective pharmacological tools. The authors describe the experimental strategies employed to investigate the poorly known role of  $\text{Ca}^{2+}$  channels in diseases associated

with neuronal migration. An interesting example, recently clarified, is Fetal Minamata disease (FMD), caused by exposure to methylmercury (MeHg) during development and associated with disrupted neuronal migration, maturation, and growth. MeHg inhibits granule cell migration by reducing the frequency of somal  $\text{Ca}^{2+}$  spikes through alterations in  $\text{Ca}^{2+}$ , cAMP, and insulin-like growth factor 1 (IGF1) signaling [37]. On the other hand, Lovisolo et al. investigate the putative roles of  $\text{Ca}^{2+}$ -dependent events that occur during postnatal physiological neurogenesis in the olfactory system. A general and debated question, critically analysed in this review, concerns the relative contribution of voltage-dependent and voltage-independent  $\text{Ca}^{2+}$  channels in the control of functional properties of neurosecretory cells and neuronal motility [38].

The review by Moccia et al. explores the debated involvement of endothelial progenitor cells (EPCs) in tumor neovascularisation and progression [39]. This event is suggested to integrate the classical angiogenic pathway due to sprouting from pre-existing vessels. Growing tumors may recruit EPCs from bone marrow to trigger blood vessel formation through physical engrafting into the vessel lumen and/or paracrine release of proangiogenic factors. Tumor cell-based therapy (CBT) is limited by the paucity of EPCs harvested from peripheral blood and suffers from several drawbacks [40]. A better understanding of cell signaling that controls EPC homing, proliferation and incorporation into injured tissues would help to improve this strategy. In particular, the authors investigate the potential clinical relevance of store-operated  $\text{Ca}^{2+}$  entry (SOCE), a  $\text{Ca}^{2+}$ -permeable membrane pathway that is activated upon depletion of intracellular  $\text{Ca}^{2+}$  pools. SOCE may actually contribute to the proangiogenic effects exerted by vascular endothelial growth factor (VEGF) in subpopulations of circulating EPCs.

The increasing amount of information on intracellular calcium dynamics clearly indicates that human diseases are often associated with disorders in  $\text{Ca}_c$  signaling. However, the assumption that ‘correlation implies causation’ should not be assumed *a priori* and warrants critical consideration. We are only beginning to integrate the plethora of experimental evidences obtained both *in vitro* and *in vivo* by the use of different models (from cell lines to animals, in healthy and altered tissues) and techniques (electrophysiology, live cell imaging). The differential structure, expression, function or targeting of  $\text{Ca}^{2+}$ -toolkit components can truly concur to the pathogenesis of some disorders, as in the case of channelopathies, in which  $\text{Ca}_c$  signals are

suitable selective targets for therapeutical approaches [11, 41-46]. Conversely, disease progression may often lead to secondary effects involving  $\text{Ca}_c$  signaling machinery. In this case, we could look at them as putative diagnostic biomarkers, mostly when they are altered in the early phases of the disease.

The main aim of this issue is to provide some critical hints on the fascinating contribution of calcium to the balance between health and disease, giving particular emphasis to the peculiarity of each biological process.


## References

- [1] Berridge MJ, Lipp P, Bootman MD. The versatility and universality of calcium signalling. *Nat Rev Mol Cell Biol* 2000 Oct 1;1(1):11-21.
- [2] Case RM, Eisner D, Gurney A, Jones O, Muallem S, Verkhratsky A. Evolution of calcium homeostasis: from birth of the first cell to an omnipresent signalling system. *Cell Calcium* 2007 Jan 1;42(4-5):345-50.
- [3] Clapham DE. Calcium signaling. *Cell* 2007 Dec 14;131(6):1047-58.
- [4] Collins SR, Meyer T. Evolutionary origins of STIM1 and STIM2 within ancient  $\text{Ca}^{2+}$  signaling systems. *Trends Cell Biol* 2011 Apr;21(4):202-11.
- [5] Munaron L. Shuffling the cards in signal transduction: Calcium, arachidonic acid and mechanosensitivity. *WJBC* 2011 Jan 1;2(4):59.
- [6] Gaudet R. TRP channels entering the structural era. *J Physiol* 2008 Aug 1;586(Pt 15):3565-75.
- [7] Venkatachalam K, Montell C. TRP channels. *Annu Rev Biochem* 2007;76:387-417.
- [8] Jentsch TJ, Hubner CA, Fuhrmann JC. Ion channels: function unravelled by dysfunction. *Nat Cell Biol* 2004 Nov;6(11):1039-47.
- [9] Boulware MJ, Marchant JS. Timing in cellular  $\text{Ca}^{2+}$  signaling. *Curr Biol* 2008 Sep 9;18(17):R769-R76.
- [10] Munaron L, Fiorio Pla A. Endothelial calcium machinery and angiogenesis: understanding physiology to interfere with pathology. *Curr Med Chem* 2009;16(35):4691-703.
- [11] Nilius B. TRP channels in disease. *Biochim Biophys Acta* 2007 Aug 1;1772(8):805-12.
- [12] Balla T. Regulation of  $\text{Ca}^{2+}$  entry by inositol lipids in mammalian cells by multiple mechanisms. *Cell Calcium* 2009 Jun;45(6):527-34.

- [13] Michell RH. First came the link between phosphoinositides and Ca<sup>2+</sup> signalling, and then a deluge of other phosphoinositide functions. *Cell Calcium* 2009 Jun;45(6):521-6.
- [14] Wang R. Physiological implications of hydrogen sulfide: a whiff exploration that blossomed. *Physiol Rev* 2012 Apr;92(2):791-896.
- [15] Hardie RC, Muallem S. Lipids in Ca<sup>2+</sup> signalling--an introduction. *Cell Calcium* 2009 Jun 1;45(6):517-20.
- [16] Mustafa AK, Gadalla MM, Snyder SH. Signaling by gasotransmitters. *Science Signaling* 2009;2(68):re2.
- [17] Halls ML, Cooper DM. Regulation by Ca<sup>2+</sup>-signaling pathways of adenylyl cyclases. *Cold Spring Harb Perspect Biol* 2011 Jan;3(1):a004143.
- [18] Willoughby D. Organization of cAMP signalling microdomains for optimal regulation by Ca<sup>2+</sup> entry. *Biochem Soc Trans* 2012 Feb;40(1):246-50.
- [19] Lefkimmatis K, Srikanthan M, Maiellaro I, Moyer MP, Curci S, Hofer AM. Store-operated cyclic AMP signalling mediated by STIM1. *Nat Cell Biol* 2009 Apr 1;11(4):433-42.
- [20] Zaccolo M. Spatial control of cAMP signalling in health and disease. *Curr Opin Pharmacol* 2011 Dec;11(6):649-55.
- [21] Siso-Nadal F, Fox JJ, Laporte SA, Hébert TE, Swain PS. Cross-talk between signaling pathways can generate robust oscillations in calcium and cAMP. *PLoS ONE* 2009 Jan 1;4(10):e7189.
- [22] Spirli C, Locatelli L, Fiorotto R, Morell CM, Fabris L, Pozzan T, et al. Altered store operated calcium entry increases cyclic 3',5'-adenosine monophosphate production and extracellular signal-regulated kinases 1 and 2 phosphorylation in polycystin-2-defective cholangiocytes. *Hepatology* 2012 Mar;55(3):856-68.
- [23] Parekh AB. Ca<sup>2+</sup> microdomains near plasma membrane Ca<sup>2+</sup> channels: impact on cell function. *J Physiol (Lond)* 2008 Jan 1.
- [24] Rizzuto R, Pozzan T. Microdomains of intracellular Ca<sup>2+</sup>: molecular determinants and functional consequences. *Physiol Rev* 2006 Jan;86(1):369-408.
- [25] Ambudkar IS. Dissection of calcium signaling events in exocrine secretion. *Neurochem Res* 2011 Jul;36(7):1212-21.
- [26] Petersen OH, Gerasimenko OV, Gerasimenko JV. Pathobiology of acute pancreatitis: focus on intracellular calcium and calmodulin. *F1000 Med Rep* 2011;3:15.
- [27] Kim MS, Lee KP, Yang D, Shin DM, Abramowitz J, Kiyonaka S, et al. Genetic and pharmacologic inhibition of the Ca<sup>2+</sup> influx channel TRPC3 protects secretory epithelia from Ca<sup>2+</sup>-dependent toxicity. *Gastroenterology* 2011 Jun;140(7):2107-15, 15 e1-4.
- [28] Fearnley CJ, Roderick HL, Bootman MD. Calcium signaling in cardiac myocytes. *Cold Spring Harb Perspect Biol* 2011 Nov;3(11):a004242.
- [29] Weiss JN, Nivala M, Garfinkel A, Qu Z. Alternans and arrhythmias: from cell to heart. *Circ Res* 2011 Jan 7;108(1):98-112.



- [30] Hausenloy DJ, Yellon DM. The therapeutic potential of ischemic conditioning: an update. *Nat Rev Cardiol* 2011 Nov;8(11):619-29.
- [31] Hausenloy DJ, Yellon DM. Preconditioning and postconditioning: united at reperfusion. *Pharmacol Ther* 2007 Nov;116(2):173-91.
- [32] Hausenloy DJ, Yellon DM. Cardioprotective growth factors. *Cardiovasc Res* 2009 Jul 15;83(2):179-94.
- [33] Schwartz RJ, Yeh ET. Weighing in on heart failure: the role of SERCA2a SUMOylation. *Circ Res* 2012 Jan 20;110(2):198-9.
- [34] Scianna M, Munaron L, Preziosi L. A multiscale hybrid approach for vasculogenesis and related potential blocking therapies. *Progress in biophysics and molecular biology* 2011 Aug;106(2):450-62.
- [35] Munaron L, Tomatis C, Fiorio Pla A. The secret marriage between calcium and tumor angiogenesis. *Technol Cancer Res Treat* 2008 Aug 1;7(4):335-9.
- [36] Zheng JQ, Poo MM. Calcium signaling in neuronal motility. *Annu Rev Cell Dev Biol* 2007;23:375-404.
- [37] Fahrion JK, Komuro Y, Li Y, Ohno N, Littner Y, Raoult E, et al. Rescue of neuronal migration deficits in a mouse model of fetal Minamata disease by increasing neuronal Ca<sup>2+</sup> spike frequency. *Proc Natl Acad Sci U S A* 2012 Mar 27;109(13):5057-62.
- [38] Ariano P, Dalmazzo S, Owsianik G, Nilius B, Lovisolo D. TRPC channels are involved in calcium-dependent migration and proliferation in immortalized GnRH neurons. *Cell Calcium* 2011 Jun;49(6):387-94.
- [39] Bussolati B, Grange C, Camussi G. Tumor exploits alternative strategies to achieve vascularization. *The FASEB journal : official publication of the Federation of American Societies for Experimental Biology* 2011 Sep;25(9):2874-82.
- [40] De Palma M, Nucera S. Circulating endothelial progenitors and tumor resistance to vascular-targeting therapies. *Cancer Discov* 2012 May;2(5):395-7.
- [41] Feske S. CRAC channelopathies. *Pflugers Arch* 2010 Jul;460(2):417-35.
- [42] Arcangeli A, Crociani O, Lastraioli E, Masi A, Pillozzi S, Becchetti A. Targeting ion channels in cancer: a novel frontier in antineoplastic therapy. *Curr Med Chem* 2009 Jan 1;16(1):66-93.
- [43] Prevarskaya N, Skryma R, Shuba Y. Calcium in tumour metastasis: new roles for known actors. *Nat Rev Cancer* 2011 Aug;11(8):609-18.
- [44] Cain SM, Snutch TP. Voltage-gated calcium channels and disease. *Biofactors* 2011 May-Jun;37(3):197-205.
- [45] Striessnig J, Bolz HJ, Koschak A. Channelopathies in Cav1.1, Cav1.3, and Cav1.4 voltage-gated L-type Ca<sup>2+</sup> channels. *Pflugers Arch* 2010 Jul;460(2):361-74.
- [46] Pietrobon D. CaV2.1 channelopathies. *Pflugers Arch* 2010 Jul;460(2):375-93.



**Figure legend.**

Scheme showing the role of calcium in the finely tuned balance between health and disease (picture taken from ancient Knossos palace, Krete, Greece).

Complex spatio-temporal calcium dynamics, tightly and specifically integrated with other intracellular pathways, contribute to the regulation of physiology, or can concur to functional alterations, during development, postnatal and adult life. Examples in the scheme refer to those described in more detail in the present issue and are not exhaustive.